




Official Journal Issued by  
Faculty of  
Veterinary Medicine

## Benha Veterinary Medical Journal

Journal homepage: <https://bvmj.journals.ekb.eg/>



Since 1990

### Original Paper

## Detection of virulence factors of *Mycoplasma* species isolated from chicken by multiplex PCR

Ashraf A. Abd-El-Tawab<sup>1</sup>, Wafaa M. M. Hassan<sup>2</sup>, Mohammed S. El-ordy<sup>3</sup>

<sup>1</sup> Bacteriology, Immunology and Mycology Department, Faculty of Veterinary Medicine, Benha University, Egypt.

<sup>2</sup> National Laboratory for Veterinary Quality Control on Poultry Production, Animal Health Research Institute, Dokki-Giza, Egypt.

<sup>3</sup> Veterinarian

### ARTICLE INFO

#### Keywords

Broiler chickens  
*Mycoplasma gallisepticum*  
*Mycoplasma synoviae*  
PCR  
Sensitivity test.

Received 09/2/2019

Accepted 04/03/2019

Available On-Line

18/07/2020

### ABSTRACT

One hundred and sixty clinically diseased broiler local chickens collected from 22 different farms belonging to Qalyubia, Dakahlia and Gharbia governorates were subjected to bacteriological examination and molecular characterization. Clinical signs were chronic respiratory disease with respiratory manifestations, lameness, loss or reduction in egg production. Bacteriological examination showed that 18.18% of the isolates were *Mycoplasma* positive and showed growth in pleuropneumonia-like organism (PPLo) agar plates, with fried egg appearance when examined by stereoscopic microscope. *Mycoplasma* colonies were tested for antimicrobial sensitivity tests against 12 antimicrobial antibiotics showed higher sensitivity to nitrofurantoin, gentamicin, norfloxacin, ciprofloxacin and neomycin. Intermediate sensitivity were recorded to ceftriaxone and cefotaxime, and antibiotic resistance was recorded to tetracyclines, lincomycin, chloramphenicol and sulphamethoxasine. Identification of the bacterial strains of the isolates was conducted by multiplex PCR using two primer pairs for *Mycoplasma synoviae* (v1hA) and *Mycoplasma gallisepticum* (mgc2). The amplicons expected sizes were 396 bp, and 300 bp for v1hA and mgc2, respectively. Only 13 out of 22 farms were positive, representing 59.09%. Moreover, the incidence rate of *M. synoviae* (v1hA) and *M. gallisepticum* (mgc2) was 22.72% and 13.63%, respectively, and 22.72% of the inspected farms showed positive results for both *Mycoplasma* strains. In conclusion, High prevalence of mixed *M. synoviae* and *M. gallisepticum* infections in poultry cause respiratory manifestations. Multiplex PCR is sensitive and specific for simultaneous detection of *M. synoviae* and *M. gallisepticum* in a single reaction.

## 1. INTRODUCTION

*Mycoplasma* is a member of *Mollicutes* class. *Mycoplasma* is recognized as the most important pathogen in poultry associated with high mortality rates, increase in carcass condemnations and drop in egg production (Levisohn and Kleven, 2000). Moreover, transient suppression of humeral and cellular immune responses, immune tolerance and auto immune diseases, as well as the massive lymphoid cell infiltration in the respiratory tract and joint tissues of infected fowls were occurred during *mycoplasma* infection (Yamamoto et al., 1990; Razin et al., 1998). *Mycoplasmas* infection is induced after the host is exposed to stress factors like vaccination, cold weather, overcrowding, feed/water restriction, temperature extremes, poor ventilation and other stress. *Mycoplasma* infection is usually associated with respiratory manifestations, high mortalities, reduced weight gain and condemnation of birds at the slaughter.

*Mycoplasma gallisepticum* strain was first isolated by Yoder (1980), while *M. synoviae* was first isolated from synovial sheath of commercial chickens by Morrow et al. (1990).

The most important *Mycoplasma* pathogens of the poultry are *Mycoplasma gallisepticum*, *Mycoplasma synoviae*, *Mycoplasma meleagridis* (only for turkeys) and *Mycoplasma iowae*. All of them causes significant economic losses (Kleven, 1997; Ley 1997; Yoder, 1991).

*Mycoplasmas* are thought to colonize in mucosal surfaces more efficiently and become more virulent by alternating the composition of their surface proteins. *Mycoplasma gallisepticum* cytoadhesin membrane surface proteins that undergo changes are represented by pmgA (hemagglutinins), mgc1, mgc2 and pvpA (Bencina et al. 2002). *Mycoplasma Synoviae* has two major surface antigens, that are encoded by a single gene, v1hA (variably expressed lipoprotein (MSPB) and the haemagglutinin (MSPA)) (Kiseok et al., 2010). Recent comparison of the *M. gallisepticum* genome with the *M. synoviae* genome revealed that a number of their genes have been transferred horizontally (Papazisi et al., 2003).

Although antimicrobials are considered very important method for treatment of clinical disease and maintaining birds' health and productivity, they have been implicated as risk factors in the dissemination and development of drug resistance (Whithear et al. 1983). *M. gallisepticum* may

\* Corresponding author: Prof. Ashraf A. Abd-El-Tawab, Bacteriology, Immunology and Mycology Department, Faculty of Veterinary Medicine, Benha University, Egypt.

develop resistance and even demonstrate cross-resistance to commonly used antibiotics (Zanella et al., 1998). Moreover, *Mycoplasma* revealed sensitivity to tetracyclines and quinolones, but they are resistant to penicillin and other antibiotic inhibitors of cell wall synthesis (Bébéar et al. 1999).

The aim of the present study was to evaluate a multiplex PCR assay for rapid detection of *Mycoplasma* pathogens in clinical specimens of chicken suffered from respiratory manifestations and loss of egg production thus would allow earlier and appropriate treatment as well as control of the diseases.

## 2. MATERIAL AND METHODS

### 2.1. Chicken samples:

A total of 160 chicken samples were collected aseptically from heart, liver, trachea and synovial fluids from 22 different farms suspected to be infected with *Mycoplasma* and belonging to Qalyubia, Dakahlia and Gharbia governorates.

### 2.2. Bacteriological examination:

Pleuropneumonia-like organism (PPO) plus DNA media (PPO broth or PPO agar, Horse serum, Yeast extract 5% solution, DNA 0.2% w/v solution, Penicillin G-Sodium and Thallium acetate 2% w/v solution) was used for bacteriological examination according to Sabry (1968).

### 2.3 Microscopical examination:

By using Geimsa staining technique as described by Sabry (1968) for morphological study.

### 2.4. Biochemical identification of the bacterial isolates:

It was performed according to Sabry (1968) including Glucose fermentation medium (Phenol red dextrose broth base, Horse serum, Thallium acetate 2% w/v solution and Penicillin G-Sodium) and Arginine deamination medium (Phenol red broth base, L-arginine solution (10% w/v solution), Horse serum, Thallium acetate 2% w/v solution and Penicillin G-Sodium)

### 2.5. Antimicrobial sensitivity test:

The disk diffusion method was applied according to Bauer et al. (1966). All *mycoplasma* isolates were tested for their antimicrobial susceptibility by 12 different antimicrobial agents (Difco™) including; neomycin (30 µg), nalidixic acid (30 µg), nitrofurantoin (300 µg), tetracycline (30 µg), ciprofloxacin (30 µg), gentamicin (10 µg), oxytetracycline (30 µg), ceftriaxone (30 µg), lincomycin (µg), norfloxacin (10 µg), cefotaxime (30 µg), sulphamethoxasine (100 µg), chloramphenicol (30 µg). The interpretation of inhibition zones of tested culture was done according to NCCLS, (2002), when the zone of inhibition had a diameter  $\geq 20$ mm, the isolate was considered sensitive to the used antibiotic.

### 2.6. Detection of virulence genes PCR method:

DNA extraction was performed using QIA amp DNA mini extraction kit according to the manufacturer's instruction. DNA was amplified by using the PCR method (Lysnyansky et al., 2005). PCR Master Mix used for cPCR is Emerald Amp GT PCR mastermix (Takara) Code No. RR310A

contains Emerald Amp GT PCR mastermix (2x premix) and PCR grade water. Temperature and time conditions of the two primers during PCR are shown in table (1) according to specific authors and Emerald Amp GT PCR mastermix (Takara) kit.

Table 1 Cycling conditions of the different primers during cPCR.

Final extension	No. of cycles	Extension	Annealing	Secondary denaturation	Primary denaturation
72°C	35	72°C	55°C	94°C	94°C
10 min.		1 min.	30 sec.	30 sec.	5 min.

For detection of the target genes (*vilA* and *mge2*). PCR products were separated by gel electrophoresis and visualized using U.V trans-illuminator

## 3. RESULTS

### 3.1. Incidence of Mycoplasma in the examined chicken farms by using conventional culture method:

Out of 160 samples taken from diseased broiler chickens from 22 different broiler farms from Qalyubia, Dakahlia and Gharbia Governorates, 4 farms showed *mycoplasma* positive, with an incidence rate of 18.18%. Isolates appeared as fried egg when examined by stereoscopic microscope, sensitive to digitonin, positive to glucose fermentation test and negative for arginine deamination test.

### 3.2. Antimicrobial sensitivity testing of Mycoplasma isolated from broiler chickens:

Results of *in-vitro* antimicrobial sensitivity testing for *Mycoplasma* isolates were demonstrated in table (3). *Mycoplasma* isolates were highly resistant to tetracyclines. Moderate sensitivity was observed against ceftriaxone and ciprofloxacin, and highly sensitive to gentamicin and nitrofurantoin.

Table 3 Antibiogram patterns for *Mycoplasma* recovered from cases of diseased broiler chickens and local breeds' chickens

Antibacterial agents	Disc content (µg)	(n=4)			
		Susceptible		Resistant	
		No.	%	No.	%
Neomycin	3	3	75	1	25
Ceftriaxone	30	2	50	2	50
Nalidixic acid	30	3	75	1	25
Ciprofloxacin	30	3	75	1	50
Tetracycline	30	0	0	4	100
Gentamicin	10	3	75	1	25
Nitrofurantoin	30	3	75	1	25
Lincomycin	10	2	50	2	50
Chloramphenicol	30	2	50	2	50
Sulphamethoxasine	100	2	50	2	50
Norfloxacin	10	3	75	1	25
Cefotaxime	30	2	50	2	50

No.: Number of positive cases. %: was calculated according to the total number of *mycoplasma* isolates (n=4)

### 3.3. Incidence of Mycoplasma species in different poultry farms by using multiplex PCR were demonstrated in table (4)

3.4 Comparison between percentage of detection of mycoplasma from clinical samples using conventional culture methods and PCR in different governorates were demonstrated in table (5)

Table 4 Incidence of Mycoplasma species in different poultry farms by using multiplex PCR

Bacterial infections	No. of infected farms (%)	Type of production	
		Local breed	broiler
<i>M. gallisepticum</i>	3 (13.63%)	3	0
<i>M. synoviae</i>	5 (22.72%)	4	1
<i>M.gallisepticum</i> and <i>M.synoviae</i>	5 (22.72%)	4	1
No growth	9(40.9%)	5	4
Total	22 (100%)	16	6

Table 5 Comparison between percentage of detection of mycoplasma from clinical samples using conventional culture methods and PCR in different governorates

Incidence of Mycoplasma farms		Conventional culture methods	PCR
Governorate	No. of farms		
Qalyubia	17	3	11
Dakahlia	3	1	2
Gharbia	2	0	0
Total	22	4	13

#### 4. DISCUSSION

*Mycoplasma* infections are of high economic importance in the poultry industry because high mortality rates, poor carcass conditions and loss of egg production. Avian *mycoplasmas* are induced after the host is affected by other disease-causing agents such as bacteria and viruses and/or after an episode of host weakness (Yoder et al., 1991). Interestingly, the classical microbiological techniques currently in use for *Mycoplasma* detection and identification are not satisfactory in most situations but remain necessary for drug susceptibility testing. The complexity associated with them makes alternative approaches more attractive (Anbazhagan et al., 2010). In our study, from 22 examined poultry farms and by conventional identifications methods, *Mycoplasma* was recorded in 4 farms (18.18 %). Yoder (1984) stated that *Mycoplasma* were fastidious. They were more sensitive than bacteria to environmental conditions such as tonicity of the medium and the exposed plasma membrane was sensitive to damage by surface active substance. This explains the decreased chances for isolation of *Mycoplasma* by conventional methods.

*M. gallisepticum* was isolated from chickens showing signs of chronic respiratory disease with a percentage of 5% (Abd El Aziz et al., 2007). Meanwhile, Hassan (2001) isolated *M. gallisepticum* with an incidence rate 14.6%. Also, Heleili et al. (2011) isolated *M. gallisepticum* from respiratory organs of chickens with a percentage of 21.67%. These results are agreed with our detection result (18.18%) which confirm low detection rates for *Mycoplasma* by conventional methods due to loss of extra *mycoplasma* cells during cultivation and cell membrane damages in comparison to PCR results.

PCR was used to assess the prevalence of microorganisms incriminated in occurrence of CRD signs in poultry, and the results were compared with those obtained using culture techniques. Obviously, the PCR assays have demonstrated a significantly higher rate of detection of *Mycoplasma* in poultry farms with various problems in Egypt than detection by classical culture procedures. Overall, PCR could detect *Mycoplasma* in 13 farms (59.09%). Also, Marois et al. (2002) recorded positive *M. gallisepticum* cases through

culture identification 3.75% in comparison to molecular technique (42.4%). Moreover, Rauf et al. (2013) used 16S rRNA gene as species specific primers of *MG* and found overall 27.6% from field birds were positive for *MG* by conventional cultivation methods in comparison to PCR (68.94%). These results are attributed to the fact that PCR can detect DNA from both viable and non-viable bacteria and hence is more reliable diagnostic test in terms of sensitivity and specificity. Thus, the use of reference genes differ from genes applied in the previous studies, the results are similar to our results and *Mycoplasma* showed different isolation results for *MG* and *MS* at the same sample.

In a previous study, Boussetta et al., (1997) isolated *MG* from 15 flocks (23.8 %), while *M.S* was isolated from only five flocks (7.9 %) in Tunisia. The prevalence of *M.S* in backyard chickens averaged between 68.6 % and 100%, while the prevalence of *M.G* was averaged between 32.8% and 55.1% (Xavier et al., 2011). But in our study, overall, 5 out of 22 examined farms were assigned to be positive for *M. synoviae* (22.72%) were with single infection, 3 (13.63%) associated with *M. gallisepticum* and 5 (22.72) showed mixed infections, meanwhile 9 farms only (40.9%) were negative.

Multiplex PCR assay was optimized for successful detection of genes with expected amplicon sizes from clinical specimens collected from suspected farms. The results showed that mPCR yielded a detectable DNA fragment of expected molecular weight only in the presence of their respective DNA template and gave negative results when tested with other bacteria.

Siddique et al. (2012) optimized the multiplex PCR for successful detection of five of the respiratory tract pathogens including *M. gallisepticum*, *M. synoviae*, Newcastle disease virus, Infectious bronchitis virus and Avian influenza virus. Bayatzadehet al. (2011) amplified the conserved region of 16S rRNA gene for the detection of *Mycoplasma* genus in 163bp fragment and *M. synoviae* in 207bp.

In the current work, two reference strains of *Mycoplasma* including *M. synoviae* (*vlhA*) and *M. gallisepticum* (*mgc2*) with expected amplicon sizes 396bp and 300bp respectively, which amplified using its respective primer pairs.

In general, the use of multiplex PCR reactions for groups of organisms causing similar syndromes provides an efficient way to ask several related epidemiological questions simultaneously. On the other hand, vaccination didn't give complete protection against infection, but some were effective in suppress the multiplication of the organism, resulting in less tissue damage followed by faster recovery (Hildebrand et al., 1983; Rodriguez and Kleven, 1985).

Antibiotic treatment (chemotherapy) is necessary in complement of biosecurity to control *Mycoplasma* infections. It is logic that for a successful and aimed *mycoplasma* infection treatment, it is necessary to have regular antibiogram tests of *M. gallisepticum* and or *M. synoviae* in the field for monitoring susceptibility of *Mycoplasma* prevalent in the farms. In previous studies, *Mycoplasma* was reported to show sensitivity *in vitro* and *in vivo* to tetracyclines and quinolones (Jordan and Horrocks 1996; Bébear et al., 1999; Wu et al., 2000) which is opposite to our results. On the other hand, our results came in agreement with that reported by Whithear et al. (1983), who recorded that *Mycoplasma* isolates showed resistance to oxytetracycline and erythromycin.

## 5. CONCLUSION

*M. synoviae* consider as the most common poultry mycoplasma causes respiratory infections in poultry. Higher prevalence of mixed *M. synoviae* and *M. gallisepticum* infections in poultry with respiratory manifestations was recorded. Multiplex PCR is sensitive and specific for simultaneous detection of *M. synoviae* and *M. gallisepticum* in clinical specimens in a single reaction. *Mycoplasma* are highly sensitive to gentamicin and nitrofurantoin antibiotics, while they are highly resistant to tetracyclines

## 6. REFERENCES

1. Abd El Aziz, E.E.; Hassan, A.M.; Badr, J.M. (2007): In vitro efficacy of some antimicrobials on the *E. coli* and Mycoplasmal isolates from cases of chronic respiratory disease in broilers in Egypt. *Zag. Vet. J.* 35 (3): 40-49.
2. Anbazhagan, D.; Mui, W.S.; Mansor M.; Yan, G.O.S.; Yusof, M.Y.; Sekaran, S.D. (2011): Development of conventional and real-time multiplex PCR assays for the detection of nosocomial pathogens. *Brazilian J. Microbiol.* 42: 448-458.
3. Bébéar, C.M.; Renaudin, J.; Charon, A.; Renaudin, H.; de Barbeyrac, B.; Schaefferbeke T.; Bébéar. C. (1999): Mutations in the *gyrA*, *parC* and *par E* genes associated with fluoroquinolones resistance in clinical isolates of *Mycoplasma hominis*. *Antimicrob. Agents Chemoter.*, 43: 954-956.
4. Bauer, A.W.; Kirby, W.M.; Sherris, J.C.; Turck. M. (1966): Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Pathol.* 45: 493-496.
5. Bayatzadeh, M.A.; Pourbakhsh, S.A.; Homayounimehr, A.R.; Ashtari, A.; Abtin, A.R. (2011) : Application of culture and polymerase chain reaction (PCR) methods for isolation and identification of *Mycoplasma synoviae* on broiler chicken farms. *Arch Razi Inst.* 66: 87-94.
6. Bencina D. (2002): Hemmagglutinins of pathogenic avian mycoplasmas. *Avian pathol.* 31:535-547.
7. Boussetta M.; Chaouachi N.; Mlik B. (1997): Etude etivitis apparently caused by *Mycoplasma gallisepticum* in sérologique et bactériologique des mycoplasmosesaviaires dans la région du Cap Bon enTunisie. *Rev.Elév.Med.Vet.* layer chickens. *Vet. Pathol.* 32: 11-18.
8. García, M.; Ikuta N.; Levisohn S.; Kleven S.H. (2005): Evaluation and comparison of various PCR methods for detection of *Mycoplasma gallisepticum* infection in chickens. *Avian Dis.* 49: 125-132.
9. Hassan, A.M. (2001): Recent techniques in diagnosis of *Mycoplasma gallisepticum* and *M. synoviae* in poultry. Thesis, Ph.D., Faculty of Veterinary Medicine, Cairo University.
10. Heleili, N.; Mamache, B.; Chelih, A. (2011): Incidence of avian mycoplasmosis in the region of Batna, Eastern Algeria. *Veterinary World J.* 4(3): 101-105.
11. Hildebrand, D.G.; Page, D.E.; Berg, J.R. (1983): *Mycoplasma gallisepticum* (MG) — laboratory and field studies evaluating the safety and efficacy of an inactivated MG bacterin. *Avian Dis.* 27: 792-802.
12. Hirose, K.; Kobayashi, H.; Ito, N.; Kawasaki, Y.; Zako, M.; Kotani, K.; Ogawa, H.; Sato, H. (2003): Isolation of *Mycoplasmas* from nasal swabs of calves affected with respiratory diseases and antimicrobial susceptibility of their isolates. *J. Vet. Med.* 50: 347-351.
13. Hong, Y.; Garcia, M.; Leiting, V.; Bencina, D.; Dufour-Zavala, L.; Zavala, G. and Kleven, S.H. (2004): Specific Detection and Typing of *Mycoplasma synoviae* Strains in Poultry with PCR and DNA Sequence Analysis Targeting the Hemagglutinin Encoding Gene *vlhA*. *Avian Diseases* 48:606-616.
14. Jordan, F.T.W.; Horrocks. B.K. (1996): The minimum inhibitory concentration of tilmicosin and tylosin for *Mycoplasma gallisepticum* and *Mycoplasma synoviae* and a comparison of their efficacy in the control of *Mycoplasma gallisepticum* in broiler chicks. *Avian Dis.* 40: 326-334.
15. Kleven, S.H. (1985): Tracheal populations of *Mycoplasma gallisepticum* after challenge of bacterin-vaccinated chickens. *Avian Diseases* 29: 1012-1017
16. Kleven, S.H. (1997): *Mycoplasma synoviae* infection in: *Diseases of poultry*, ed, Calnek BW, 10th ed., pp.220-225. Iowa state university Press, Ames, IA.
17. Levisohn, S.; Kleven, S.H. (2000): Avian mycoplasmosis (*Mycoplasma gallisepticum*). *Rev. Sci. Tech.* 19(2): 425-442.
18. Ley, D.H.; Yoder, H.W.Jr. (1997): *Mycoplasma gallisepticum*. In *Diseases of Poultry*. 10th ed. Calnek, B. W. (pp. 194-207): Ames, IA: Iowa state University press.
19. Lysyansky, I.; Garcia, M.; Levisohn, S. (2005): Use of *mgc2*-Polymerase chain reaction-Restriction fragment length polymorphism for rapid differentiation between field isolates and vaccine strains of *Mycoplasma*. *Avian Dis.* 49(2):238-45
20. Marios, C.; Gebert, F. D.; Kempf, I. (2002): Polymerase chain reaction for detection of *Mycoplasma gallisepticum* in environmental sample. *Avian Pathol.* 31: 163-168.
21. Morrow, C.J.; Bell, I.G.; Walker, S.B.; Markham, P.F.; Thorp, B.H.; Whithear, K.G. (1990): Isolation of *Mycoplasma synoviae* from infectious synovitis of chickens. *Australian Vet. J.* 66(4), 121-124.
22. NCCLS. (2002): Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated from Animals: Approved Standard M31-A2. 2nd Edn., National Committee for Clinical Laboratory Standards, Wayne, PA., USA.
23. OIE-World Organisation for Animal Health. (2010): Avian mycoplasmosis (*Mycoplasma gallisepticum*), In, Anna RovidSpickler, Roth, A.J, Galyon, J. and Lofstedt, J. (Eds.): *Emerging and Exotic Diseases of Animals*, Iowa State University, USA: 94-97.
24. Oh, K.; Lee, S.; Seo, J.; Lee, D.; Kim, T. (2010): Rapid serodiagnosis with the use of surface plasmon resonance imaging for the detection of antibodies against major surface protein A of *Mycoplasma synoviae* in chickens. *Can. J. Vet. Res.* 74(1):71-74.
25. Papazisi, L.; Gorton, T.S.; Markham, P.F.; Browning, G.F.; Ngguyen, D.K.; Swartzell, S.; Madan, A.; Mahairas, G.; Geary, J.S. (2003): The complete genome sequence of the avian pathogen *Mycoplasma gallisepticum* strain Rlow. *Microbiology* 2003; 149:2307-2316.
26. Rauf, M.; Chaudhary, Z.I.; Younus, M.; Anjum, A.A.; Ali, M.A.; Ahmad, A.N.; Khan M.U.R. (2013): Identification of *Mycoplasma gallisepticum* by polymerase chain reaction and conventional diagnostics from white leghorn layer flocks. *The J. Anim. Plant Sci.* 23 (2): 393-397.
27. Razin, S.; Yogev, D.; Naot, Y. (1998): Molecular biology and pathogenicity of *Mycoplasmas*. *Microbiol. Mol. Biol. Rev.*, 62 (4): 1094-1156.
28. Sabry, M.Z. (1968): Characterization and classification of avian mycoplasmas. Thesis, Ph.D. Cornell University, USA.
29. Sabry, M.Z.; Ahmed, A.A. (1975): Evaluation of culture procedure for primary isolation of *Mycoplasmas* from female genitalia of farm animals. *J. Egypt. Vet. Med. Ass.* 35: 18-34.
30. Sabry, M.Z.; Erno, H.; Freundt, E.A. (1971): Manual of technical methods for the characterization and serotyping of mycoplasmas. *Mycoplasma Research Department, Animal Health Research Institute, Ministry of Agriculture, Cairo, Egypt.*
31. Siddique, A.B.; Rahman, S.U.; Hussain, I.; Muhammad, G. (2012): Frequency distribution of opportunistic avian pathogens in respiratory distress Cases of poultry. *Pak. Vet. J.* 32(3):386-389.
32. Vandekerchove, D.; De Herdt, P.; Laevens, H.; Butaye, P.; Meulemans, G.; Pasmans, F. (2004): Significance of interactions between *Escherichia coli* and respiratory

- pathogens in layer hen flocks suffering from colibacillosis-associated mortality *Avian Pathol.* 33(3):298-302.
33. Whithear, K.G.; Bowtell, D.D.; Ghiocas E.; Hughes, K.L. (1983): Evaluation and use of a Micro-broth dilution procedure for testing sensitivity. *Avian Dis.* 27: 937-942.
  34. Wu, C.C.; Shryock, T.R.; Lin, T.L.; Faderan. M. (2000): Antimicrobial susceptibility of *Mycoplasma hyorhinis*. *Vet. Microbiol.* 76: 25-30.
  35. Xavier, J.; Pascal, D.; Crespo, E.; Schell, H.; Trinidad, J.; Bueno, D. (2011): Seroprevalence of salmonella and mycoplasma infection in backyard chickens in the state of Entre Ríos in Argentina. *Poult. Sci. J.* 90:746-751.
  36. Yamamoto, R. (1990): Mollicutes. In: Biberstein EL, Zee YC, editors. Review of veterinary microbiology. Chicago:Blackwell Scientific Publications; p.213-27.
  37. Yoder, Jr.H.W. (1984): *Mycoplasma gallisepticum* infection. In Diseases of poultry. 8th Ed. Edited by Calnek, B. W.; Beard, C. W.; Barnes, H. J.; Reid, W. M. and Yoder Jr. H. W. Iowa State University Press, Ames, Iowa.
  38. Yoder, Jr. H. W.(1991): *Mycoplasma gallisepticum* infection. In: Calnek BW, Burnes HJ, Beard CW, Yoder Jr. HW, editors. (1999): Diseases of poultry. Ames, Iowa, USA. Ames: Iowa State University Press.Pp. 198-212.
  39. Zanella, A., P.A. Martino, A. Pratelli and M. Stonfer. (1998): Development of antibiotic resistance in *Mycoplasma gallisepticum* in vitro. *Avian Pathol.* 27: 591-596.